

### **Axillopexy**

V. Sârbu<sup>1</sup>, R. Melihov<sup>2</sup>, O. Unc<sup>2</sup>, D. Costea<sup>2</sup>, M. Pundiche<sup>2</sup>, R. Ilie<sup>3</sup>, V. Truta<sup>3</sup>

<sup>1</sup>Ovidius University, "Medstar 2000" General Surgery Clinic, Constanța, Romania

<sup>2</sup>Ovidius University, General Surgery Department of the Emergency Clinical County Hospital  
Constanța, Romania

<sup>3</sup>Municipal Hospital Mangalia, Romania

### **Abstract**

The present paper is a presentation of our technique of axillopexy, used after the excision of the axillary lymph nodes in 29 cases of breast cancer patients. We have used this technique after Madden modified radical mastectomy or after quadrantectomy for tumors in the external quadrants of the mammary gland. We have studied and compared with a 30 case control group, the duration of the lymphorrhagia the moment of removing the drains, the presence/absence of other local complications. We have also measured the time until the beginning of the oncological postoperative therapy. Every one of the aspects we have studied was improved in the axillopexy group.

**Key words:** axillopexy, limphoragy, breast cancer

Corresponding author: Prof. Univ. Dr. Vasile Sârbu  
Ovidius University Constanța, Romania  
E-mail: vasilesarbu\_cta@yahoo.com